



PubMed Nucleotide Protein Genome Structure PopSet Taxonomy OMIM Books

Search PubMed for

Go Clear

Limits Preview/Index History Clipboard Details

About Entrez

Display Abstract Sort Save Text Clip Add Order

Text Version

☐ 1: J Clin Invest 1993 Jun;91(6):2850-6

Related Articles, Links

Entrez PubMed

[Overview](#)
[Help | FAQ](#)
[Tutorial](#)
[New/Noteworthy](#)
[E-Utilities](#)[PubMed Services](#)
[Journal Browser](#)
[MeSH Browser](#)
[Single Citation Matcher](#)
[Batch Citation Matcher](#)
[Clinical Queries](#)
[LinkOut](#)
[Cubby](#)[Related Resources](#)
[Order Documents](#)
[NLM Gateway](#)
[TOXNET](#)
[Consumer Health](#)
[Clinical Alerts](#)
[ClinicalTrials.gov](#)
[PubMed Central](#)[Privacy Policy](#)

Tissue factor pathway inhibitor reduces mortality from Escherichia coli septic shock.

Creasey AA, Chang AC, Feigen L, Wun TC, Taylor FB Jr, Hinshaw LB.


Chiron Corporation, Emeryville, California 94608.

This study was designed to test the hypothesis that tissue factor pathway inhibitor (TFPI) plays a significant role in vivo in regulating coagulation that results from exposure of blood to tissue factor after vascular injury as in the case of gram negative sepsis. Highly purified recombinant TFPI (6 mg/kg) was administered either 30 min or 4 h after the start of a lethal intravenous Escherichia coli infusion in baboons. Early posttreatment of TFPI resulted in (a) permanent seven-day survivors (5/5) with significant improvement in quality of life, while the mean survival time for the controls (5/5) was 39.9 h (no survivors); and (b) significant attenuations of the coagulation response and various measures of cell injury, with significant reductions in pathology observed in E. coli sepsis target organs, including kidneys, adrenals, and lungs. TFPI administration did not affect the reduction in mean systemic arterial pressure, the increases in respiration and heart rate, or temperature changes associated with the bacterial infusion. TFPI treated E. coli infected baboons had significantly lower IL-6 levels than their phosphate buffered saline-treated controls, however tumor necrosis factor levels were similarly elevated in both groups. In contrast to the earlier 30-min treatment, the administration of TFPI at 4 h, i.e., 240 min, after the start of bacterial infusion resulted in prolongation of survival time, with 40% survival rate (2/5) and some attenuation of the coagulopathic response, especially in animals in which fibrinogen levels were above 10% of normal at the time of TFPI administration. Results provide evidence for the significance of tissue factor and tissue factor pathway inhibitor in bacterial sepsis, and suggest a role for blood coagulation in the regulation of the inflammatory response.

PMID: 8514893 [PubMed - indexed for MEDLINE]

Display Abstract Sort Save Text Clip Add Order

[Write to the Help Desk](#)[NCBI | NLM | NIH](#)[Department of Health & Human Services](#)[Freedom of Information Act | Disclaimer](#)



PubMed Nucleotide Protein Genome Structure PopSet Taxonomy OMIM Books

Search PubMed for Go Clear

Limits Preview/Index History Clipboard Details

About Entrez

Display Abstract Sort Save Text Clip Add Order

Text Version

☐ 1: Gene 1993 Dec 27;137(1):25-31

Related Articles, Links

Entrez PubMed

[Overview](#)
[Help | FAQ](#)
[Tutorial](#)
[New/Noteworthy](#)
[E-Utilities](#)

PubMed Services

[Journal Browser](#)
[MeSH Browser](#)
[Single Citation Matcher](#)
[Batch Citation Matcher](#)
[Clinical Queries](#)
[LinkOut](#)
[Cubby](#)

Related Resources

[Order Documents](#)
[NLM Gateway](#)
[TOXNET](#)
[Consumer Health](#)
[Clinical Alerts](#)
[ClinicalTrials.gov](#)
[PubMed Central](#)[Privacy Policy](#)

The discovery and characterization of a novel nucleotide-based thrombin inhibitor.

Griffin LC, Toole JJ, Leung LL.

Gilead Sciences, Foster City, CA 94404.


Thrombin is a serine protease that plays a pivotal role in thrombosis and hemostasis, and is a major target for anticoagulation and cardiovascular disease therapy. Using a novel in vitro selection/amplification technique, we have identified a new class of thrombin inhibitors based on single-stranded DNA (ssDNA) oligodeoxyribonucleotides (oligos). These thrombin inhibitors are the first example of the use of this technique to obtain ssDNA oligos that bind a target protein that does not interact physiologically with nucleic acid. Here, we review how iterative selection and amplification were used to identify short ssDNA sequences that bind and inhibit thrombin (Bock et al., *Nature* 355 (1992) 564-566), and the tertiary structure of one aptamer sequence (Wang et al., *Biochemistry* 32 (1993) 1899-1904). Results from in vitro and in vivo studies are also summarized (Griffin et al., *Blood* 81 (1993) 3271-3276). The discovery of a new class of thrombin inhibitors using this technology demonstrates the power of this new approach for rapid drug discovery and development.

PMID: 8282198 [PubMed - indexed for MEDLINE]

Display Abstract Sort Save Text Clip Add Order

[Write to the Help Desk](#)[NCBI](#) | [NLM](#) | [NIH](#)[Department of Health & Human Services](#)[Freedom of Information Act](#) | [Disclaimer](#)

i686-pc-linux-gnu Aug 30 2002 15:17:13



PubMed Nucleotide Protein Genome Structure PopSet Taxonomy OMIM Books

Search PubMed

☐ for

Go

Clear

Limits

Preview/Index

History

Clipboard

Details

About Entrez

Display

Abstract

☐ Sort

☐ Save

Text

Clip Add

Order

Text Version

☐ 1: Br J Haematol 1993 Dec;85(4):761-8

Related Articles, [NEW](#) Links

Entrez PubMed

Overview

Help | FAQ

Tutorial

New/Noteworthy

E-Utilities

PubMed Services

Journal Browser

MeSH Browser

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

LinkOut

Cubby

Related Resources

Order Documents

NLM Gateway

TOXNET

Consumer Health

Clinical Alerts

ClinicalTrials.gov

PubMed Central

Privacy Policy

Autoantibodies directed against the epidermal growth factor-like domains of thrombomodulin inhibit protein C activation in vitro.

Oosting JD, Preissner KT, Derksen RH, de Groot PG.

Department of Haematology, University Hospital Utrecht, The Netherlands.

No consensus has been obtained about the question whether autoantibodies, in particular antiphospholipid antibodies (aPL), may cause thrombosis by inhibiting thrombomodulin (TM) mediated protein C activation. In order to clarify the mechanism by which autoantibodies inhibit TM-mediated protein C activation, we have screened 12 patients with autoimmune diseases for the presence of circulating autoantibodies inhibiting TM function. In a cross-sectional study we found that IgG fractions from two patients (who were aPL negative) inhibited TM mediated protein C activation in an assay system using purified components. A longitudinal study of six patients with a history of thrombosis of which two were aPL positive showed that all had at some time circulating antibodies inhibiting TM function, suggesting that the presence of these antibodies is transient. Three different TMs were used to identify the epitope of the antithrombomodulin antibodies (aTM): rabbit TM, which contains the entire TM molecule; Solulin, which contains the extracellular part of TM, and rEGF-TM, which contains the six epidermal growth factor (EGF) domains of TM. We showed that the aTM inhibited protein C activation mediated by all three TMs, indicating that the aTM are directed against the region containing the EGF domains. When TM was incorporated in phospholipid vesicles, no inhibition by these aTM could be demonstrated. In addition, protein C activation mediated by cultured endothelial cells (EC) could not be inhibited by aTM. The lack of inhibition of TM in phospholipid vesicles and EC-TM by a TM suggests that aTM only inhibit soluble TM. In conclusion, we demonstrated the transient presence of circulating autoantibodies directed against the region of TM containing the EGF domains in SLE patients with a history of thrombotic complications. We postulate that the presence of antibodies to soluble TM may be, in addition to aPL, a risk factor for the occurrence of thrombosis in patients with autoimmune diseases.

PMID: 7522520 [PubMed - indexed for MEDLINE]

Display

Abstract

☐ Sort

☐ Save

Text

Clip Add

Order

[Write to the Help Desk](#)



PubMed Nucleotide Protein Genome Structure PopSet Taxonomy OMIM Books

Search for

Limits Preview/Index History Clipboard Details

About Entrez

Text Version

☐ 1: Thromb Haemost 1993 Sep 1;70(3):448-53

Related Articles, Links

Entrez PubMed

Overview
Help | FAQ
Tutorial
New/Noteworthy
E-Utilities

PubMed Services
Journal Browser
MeSH Browser
Single Citation Matcher
Batch Citation Matcher
Clinical Queries
LinkOut
Cubby

Related Resources
Order Documents
NLM Gateway
TOXNET
Consumer Health
Clinical Alerts
ClinicalTrials.gov
PubMed Central

Privacy Policy

The significance of TFPI in clotting assays--comparison and combination with other anticoagulants.

Nordfang O, Kristensen HI, Valentin S, Ostergaard P, Wadt J.

Novo Nordisk A/S, Gentofte, Denmark.

The anticoagulant activities of Tissue Factor Pathway Inhibitor (TFPI), heparin and hirudin were compared in intrinsic (APTT) and extrinsic (PT) activated clotting assays. In contrast to the thrombin inhibitor hirudin, heparin was 10 fold more potent in the APTT assay than in the PT assay, indicating that inhibition of intrinsic activation is important for the anticoagulant activity of heparin as measured in an APTT assay. TFPI was most potent in the PT assay and the effect of TFPI was most pronounced in the presence of other anticoagulants (heparin and hirudin). The activities of the two natural anticoagulants antithrombin III (ATIII) and TFPI were compared in a PT assay with very dilute tissue factor. In this assay system TFPI in normal plasma affected the clotting time more than ATIII in the plasma. However, when heparin was added ATIII was the major anticoagulant, but profound prolongation of the clotting time was only seen when TFPI was also added. In an ATIII deficient plasma heparin did not augment the effect of TFPI, showing that the increased effect of TFPI in the presence of heparin is dependent on the anticoagulant activity of ATIII/heparin. The effect of TFPI at prolonged clotting times was also illustrated by the significant effect of blocking TFPI in the plasma from warfarin-treated patients. Thus TFPI is a major anticoagulant in normal plasma and the effect of TFPI is especially seen at prolonged clotting times.

PMID: 8259547 [PubMed - indexed for MEDLINE]

[Write to the Help Desk](#)

[NCBI](#) | [NLM](#) | [NIH](#)

[Department of Health & Human Services](#)

[Freedom of Information Act](#) | [Disclaimer](#)

i686-pc-linux-gnu Aug 30 2002 15:17:13